

**ON SINGULARITIES OF THE INVERSE PROBLEMS
ASSOCIATED WITH PERFECT CUBOIDS.**

JOHN RAMSDEN, RUSLAN SHARIPOV

ABSTRACT. Two cubic equations and three auxiliary equations for edges and face diagonals of a rational perfect cuboid have been recently derived. They constitute a background for two inverse problems. The coefficients of cubic equations and the right hand sides of auxiliary equations are rational functions of two rational parameters, i. e. they have denominators. Hence the inverse problems have singular points. These singular points are studied in the present paper.

1. INTRODUCTION.

A rational perfect cuboid is a rectangular parallelepiped whose edges and face diagonals are rational numbers and whose space diagonal is equal to unity: $L = 1$. Finding such a cuboid is equivalent to finding a cuboid with all integer edges and diagonals, which is an unsolved problem for many years (see [1–44]).

Let x_1, x_2, x_3 be edges of a cuboid and let d_1, d_2, d_3 be its face diagonals. Then x_1, x_2, x_3 are roots of the cubic equation

$$x^3 - E_{10} x^2 + E_{20} x - E_{30} = 0. \quad (1.1)$$

Similarly, d_1, d_2, d_3 are roots of the other cubic equation

$$d^3 - E_{01} d^2 + E_{02} d - E_{03} = 0. \quad (1.2)$$

Apart from (1.1) and (1.2), the rational numbers x_1, x_2, x_3 and d_1, d_2, d_3 should obey the following three auxiliary equations:

$$\begin{aligned} x_1 x_2 d_3 + x_2 x_3 d_1 + x_3 x_1 d_2 &= E_{21}, \\ x_1 d_2 + d_1 x_2 + x_2 d_3 + d_2 x_3 + x_3 d_1 + d_3 x_1 &= E_{11}, \\ x_1 d_2 d_3 + x_2 d_3 d_1 + x_3 d_1 d_2 &= E_{12}. \end{aligned} \quad (1.3)$$

The cubic equations (1.1), (1.2) and the auxiliary equations (1.3) were obtained as a result of the series of papers [45–50]). The coefficients $E_{10}, E_{20}, E_{30}, E_{01}, E_{02}, E_{03}$ in (1.1) and (1.2) as well as the right hand sides E_{21}, E_{11}, E_{12} in (1.3) are given by explicit formulas. Here is the formula for E_{11} :

$$E_{11} = -\frac{b(c^2 + 2 - 4c)}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}. \quad (1.4)$$

2000 *Mathematics Subject Classification.* 11D25, 11D72, 14H50, 14E05.

The formulas for E_{10} , E_{01} are similar to the formula (1.4) for E_{11} :

$$E_{10} = -\frac{b^2 c^2 + 2b^2 - 3b^2 c - c}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}, \quad (1.5)$$

$$E_{01} = -\frac{b(c^2 + 2 - 2c)}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}. \quad (1.6)$$

Below are the formulas for E_{20} , E_{02} , E_{30} , E_{03} , E_{21} , E_{12} in (1.1), (1.2), and (1.3):

$$E_{20} = \frac{b}{2} (b c^2 - 2c - 2b) (2b c^2 - c^2 - 6bc + 2 + 4b) \times \\ \times (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}, \quad (1.7)$$

$$E_{02} = \frac{1}{2} (28b^2 c^2 - 16b^2 c - 2c^2 - 4b^2 - b^2 c^4 + 4b^3 c^4 - 12b^3 c^3 + \\ + 4b c^3 + 24b^3 c - 8bc - 2b^4 c^4 + 12b^4 c^3 - 26b^4 c^2 - 8b^2 c^3 + \\ + 24b^4 c - 16b^3 - 8b^4) (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}, \quad (1.8)$$

$$E_{30} = c b^2 (1 - c) (c - 2) (b c^2 - 4bc + 2 + 4b) (2b c^2 - c^2 - 4bc + \\ + 2b) (b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - 12b^2 c + 4b^2 + c^2)^{-1} \times \\ \times (bc - 1 - b)^{-2} (-c + bc - 2b)^{-2}, \quad (1.9)$$

$$E_{03} = \frac{b}{2} (b^2 c^4 - 5b^2 c^3 + 10b^2 c^2 - 10b^2 c + 4b^2 + 2bc + 2c^2 - \\ - b c^3) (2b^2 c^4 - 12b^2 c^3 + 26b^2 c^2 - 24b^2 c + 8b^2 - c^4 b + 3b c^3 - \\ - 6bc + 4b + c^3 - 2c^2 + 2c) (b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ - 12b^2 c + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (-c + bc - 2b)^{-2}, \quad (1.10)$$

$$E_{21} = \frac{b}{2} (5c^6 b - 2c^6 b^2 + 52c^5 b^2 - 16c^5 b - 2c^7 b^2 + 2b^4 c^8 - \\ - 26b^4 c^7 - 426b^4 c^5 - 61b^3 c^6 + 100b^3 c^5 + 14c^7 b^3 - c^8 b^3 - 20b c^2 - \\ - 8b^2 c^2 - 16b^2 c - 128b^2 c^4 - 200b^3 c^3 + 244b^3 c^2 + 32b c^3 + \\ + 768b^4 c^4 - 852b^4 c^3 + 568b^4 c^2 + 104b^2 c^3 - 208b^4 c + 8c^4 + \\ + 16b^3 - 112b^3 c + 142b^4 c^6 + 32b^4 - 2c^5) (b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ - 12b^2 c - 4c^3 + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}, \quad (1.11)$$

$$E_{12} = (16b^6 + 32b^5 - 6c^5 b^2 + 2c^5 b - 62b^5 c^6 + 62b^6 c^6 + 16b^4 - \\ - 180b^6 c^5 - c^7 b^3 + 18b^5 c^7 - 12b^6 c^7 - 2b^5 c^8 + b^6 c^8 + 248b^5 c^2 + \\ + 248b^6 c^2 - 96b^6 c + 321b^6 c^4 - 180b^5 c^3 - 144b^5 c - 360b^6 c^3 + \\ + b^4 c^8 + 8b^4 c^6 - 6b^4 c^7 + 18b^4 c^5 + 7b^3 c^6 + 90b^5 c^5 - 14b^3 c^5 + \\ + 17b^2 c^4 + 32b^4 c^2 + 28b^3 c^3 - 28b^3 c^2 - 4b c^3 + 8b^3 c - 57b^4 c^4 + \\ + 36b^4 c^3 - 12b^2 c^3 - 48b^4 c - c^4) (b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ - 12b^2 c + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}. \quad (1.12)$$

Based on the equations (1.1), (1.2), (1.3) and on the formulas (1.4), (1.5), (1.6), (1.7), (1.8), (1.9), (1.10), (1.11), (1.12), in [50] the following two inverse problems were formulated.

Problem 1.1. *Find all pairs of rational numbers b and c for which the cubic equations (1.1) and (1.2) with the coefficients given by the formulas (1.5), (1.7), (1.9), (1.6), (1.8), (1.10) possess positive rational roots $x_1, x_2, x_3, d_1, d_2, d_3$ obeying the auxiliary polynomial equations (1.3) whose right hand sides are given by the formulas (1.4), (1.11), (1.12).*

Problem 1.2. *Find at least one pair of rational numbers b and c for which the cubic equations (1.1) and (1.2) with the coefficients given by the formulas (1.5), (1.7), (1.9), (1.6), (1.8), (1.10) possess positive rational roots $x_1, x_2, x_3, d_1, d_2, d_3$ obeying the auxiliary polynomial equations (1.3) whose right hand sides are given by the formulas (1.4), (1.11), (1.12).*

The formulas (1.4) through (1.12) possess denominators. Therefore the inverse problems are singular for some values of b and c . The main goal of the present paper is to study these singularities.

2. THE COMMON DENOMINATOR AND ITS REDUCTION.

Let's calculate the common denominator of the fractions (1.4) through (1.12):

$$(b^2 c^4 - 6 b^2 c^3 + 13 b^2 c^2 - 12 b^2 c + 4 b^2 + c^2) (b c - 1 - b)^2 \times \\ \times (b c - c - 2 b)^2 (b^2 c^2 + 2 b^2 - 3 b^2 c + c - b c^2 + 2 b) = 0. \quad (2.1)$$

The vanishing condition (2.1) determines all singular points of the inverse problems 1.1 and 1.2. The last multiplicand in the left hand side of the formula (2.1) is taken from the denominators of (1.4), (1.5), and (1.6). Studying this multiplicand we find that it is factorable. It factors as follows:

$$b^2 c^2 + 2 b^2 - 3 b^2 c + c - b c^2 + 2 b = (b c - 1 - b) (b c - c - 2 b).$$

Applying this formula to (2.1), we reduce the equality (2.1) to

$$(b^2 c^4 - 6 b^2 c^3 + 13 b^2 c^2 - 12 b^2 c + 4 b^2 + c^2) \times \\ \times (b c - 1 - b)^3 (b c - c - 2 b)^3 = 0. \quad (2.2)$$

Some multiplicands in (2.1) are raised to the third power. But the exponents do not really affect the vanishing condition (2.2). Therefore we reduce it to

$$(b^2 c^4 - 6 b^2 c^3 + 13 b^2 c^2 - 12 b^2 c + 4 b^2 + c^2) \times \\ \times (b c - 1 - b) (b c - c - 2 b) = 0. \quad (2.3)$$

The left hand side of the formula (2.3) is the reduced common denominator of the fractions (1.4) through (1.12). It is broken into the product of three terms. Therefore singular points of the inverse problems 1.1 and 1.2 are subdivided into three singularity subvarieties.

3. THE FIRST SINGULARITY SUBVARIETY.

The first subvariety of singular points is the most simple. It is determined by the most simple factor in (2.2) through the following equation:

$$bc - 1 - b = 0. \quad (3.1)$$

The equation (3.1) is linear with respect to both b and c . Resolving it for b , we get

$$b = \frac{1}{c-1}, \text{ where } c \neq 1. \quad (3.2)$$

The formula (3.2) means that the first subvariety of singular points of the inverse problems 1.1 and 1.2 is a rational curve birationally equivalent to a straight line.

4. THE SECOND SINGULARITY SUBVARIETY.

The second subvariety of singular points is similar to the first one and is also very simple. It is determined by the following equation:

$$bc - c - 2b = 0. \quad (4.1)$$

Like in (3.2), resolving the equation (4.1) with respect to b , we get

$$b = \frac{c}{c-2}, \text{ where } c \neq 2. \quad (4.2)$$

The formula (4.2) means that the second subvariety of singular points of the inverse problems 1.1 and 1.2 is a rational curve birationally equivalent to a straight line.

5. THE THIRD SINGULARITY SUBVARIETY.

The third subvariety of singular points is different from the first two. It is determined by the following equation which is quadratic with respect to b :

$$b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - 12b^2 c + 4b^2 + c^2 = 0. \quad (5.1)$$

The discriminant of the quadratic equation (5.1) with respect to b is

$$D = -4(c-1)^2(c-2)^2 c^2. \quad (5.2)$$

Looking at the discriminant formula (5.2), we see that we can expect to find rational solutions only in one of the following three cases:

$$c = 0, \quad c = 1, \quad c = 2. \quad (5.3)$$

Substituting $c = 0$ into the formula (5.1) we derive $4b^2 = 0$. This yields one rational point at the origin of the (b, c) plane:

$$b = 0, \quad c = 0. \quad (5.4)$$

Substituting $c = 1$ and $c = 2$ into (5.1), we get two equalities $1 = 0$ and $4 = 0$ which are contradictory. Thus two of the three options in (5.3) do not actually produce any rational solutions for the equation (5.1). This result is not surprising since the equation (5.1) turns out to be reducible to $(c - 1)^2 (c - 2)^2 b^2 + c^2 = 0$.

6. CONCLUSIONS.

Despite being a fourth order equation in c , the equation (5.1) does not produce elliptic curves. Therefore the structure of the set of singular points of the inverse problems 1.1 and 1.2 is very simple. It comprises one isolated point (5.4) and two parametric subsets given by the formulas (3.2) and (4.2). This fact can be useful in computerized search for a solution of at least one of the problems 1.1 and 1.2, or in solving both of them if somehow it will be proved that the number of perfect cuboids is finite and they are in a certain range.

REFERENCES

1. *Euler brick*, Wikipedia, Wikimedia Foundation Inc., San Francisco, USA.
2. Halcke P., *Deliciae mathematicae oder mathematisches Sinnen-Confect*, N. Sauer, Hamburg, Germany, 1719.
3. Saunderson N., *Elements of algebra*, Vol. 2, Cambridge Univ. Press, Cambridge, 1740.
4. Euler L., *Vollständige Anleitung zur Algebra*, 3 Theile, Kaiserliche Akademie der Wissenschaften, St. Petersburg, 1770-1771.
5. Pocklington H. C., *Some Diophantine impossibilities*, Proc. Cambridge Phil. Soc. **17** (1912), 108–121.
6. Dickson L. E., *History of the theory of numbers*, Vol. 2: *Diophantine analysis*, Dover, New York, 2005.
7. Kraitchik M., *On certain rational cuboids*, Scripta Math. **11** (1945), 317–326.
8. Kraitchik M., *Théorie des Nombres*, Tome 3, *Analyse Diophantine et application aux cuboïdes rationnelles*, Gauthier-Villars, Paris, 1947.
9. Kraitchik M., *Sur les cuboïdes rationnelles*, Proc. Int. Congr. Math. **2** (1954), Amsterdam, 33–34.
10. Bromhead T. B., *On square sums of squares*, Math. Gazette **44** (1960), no. 349, 219–220.
11. Lal M., Blundon W. J., *Solutions of the Diophantine equations $x^2 + y^2 = l^2$, $y^2 + z^2 = m^2$, $z^2 + x^2 = n^2$* , Math. Comp. **20** (1966), 144–147.
12. Spohn W. G., *On the integral cuboid*, Amer. Math. Monthly **79** (1972), no. 1, 57–59.
13. Spohn W. G., *On the derived cuboid*, Canad. Math. Bull. **17** (1974), no. 4, 575–577.
14. Chein E. Z., *On the derived cuboid of an Eulerian triple*, Canad. Math. Bull. **20** (1977), no. 4, 509–510.
15. Leech J., *The rational cuboid revisited*, Amer. Math. Monthly **84** (1977), no. 7, 518–533; see also Erratum, Amer. Math. Monthly **85** (1978), 472.
16. Leech J., *Five tables relating to rational cuboids*, Math. Comp. **32** (1978), 657–659.
17. Spohn W. G., *Table of integral cuboids and their generators*, Math. Comp. **33** (1979), 428–429.
18. Lagrange J., *Sur le dérivé du cuboïde Eulérien*, Canad. Math. Bull. **22** (1979), no. 2, 239–241.
19. Leech J., *A remark on rational cuboids*, Canad. Math. Bull. **24** (1981), no. 3, 377–378.
20. Korec I., *Nonexistence of small perfect rational cuboid*, Acta Math. Univ. Comen. **42/43** (1983), 73–86.
21. Korec I., *Nonexistence of small perfect rational cuboid II*, Acta Math. Univ. Comen. **44/45** (1984), 39–48.
22. Wells D. G., *The Penguin dictionary of curious and interesting numbers*, Penguin publishers, London, 1986.
23. Bremner A., Guy R. K., *A dozen difficult Diophantine dilemmas*, Amer. Math. Monthly **95** (1988), no. 1, 31–36.
24. Bremner A., *The rational cuboid and a quartic surface*, Rocky Mountain J. Math. **18** (1988), no. 1, 105–121.

25. Colman W. J. A., *On certain semiperfect cuboids*, Fibonacci Quart. **26** (1988), no. 1, 54–57; see also *Some observations on the classical cuboid and its parametric solutions*, Fibonacci Quart. **26** (1988), no. 4, 338–343.
26. Korec I., *Lower bounds for perfect rational cuboids*, Math. Slovaca **42** (1992), no. 5, 565–582.
27. Guy R. K., *Is there a perfect cuboid? Four squares whose sums in pairs are square. Four squares whose differences are square*, Unsolved Problems in Number Theory, 2nd ed., Springer-Verlag, New York, 1994, pp. 173–181.
28. Rathbun R. L., Granlund T., *The integer cuboid table with body, edge, and face type of solutions*, Math. Comp. **62** (1994), 441–442.
29. Van Luijk R., *On perfect cuboids*, Doctoraalscriptie, Mathematisch Instituut, Universiteit Utrecht, Utrecht, 2000.
30. Rathbun R. L., Granlund T., *The classical rational cuboid table of Maurice Kraitchik*, Math. Comp. **62** (1994), 442–443.
31. Peterson B. E., Jordan J. H., *Integer hexahedra equivalent to perfect boxes*, Amer. Math. Monthly **102** (1995), no. 1, 41–45.
32. Rathbun R. L., *The rational cuboid table of Maurice Kraitchik*, e-print [math.HO/0111229](http://arXiv.org/math.HO/0111229) in Electronic Archive <http://arXiv.org>.
33. Hartshorne R., Van Luijk R., *Non-Euclidean Pythagorean triples, a problem of Euler, and rational points on $K3$ surfaces*, e-print [math.NT/0606700](http://arXiv.org/math.NT/0606700) in Electronic Archive <http://arXiv.org>.
34. Waldschmidt M., *Open diophantine problems*, e-print [math.NT/0312440](http://arXiv.org/math.NT/0312440) in Electronic Archive <http://arXiv.org>.
35. Ionascu E. J., Luca F., Stanica P., *Heron triangles with two fixed sides*, e-print [math.NT/0608185](http://arXiv.org/math.NT/0608185) in Electronic Archive <http://arXiv.org>.
36. Ortan A., Quenneville-Belair V., *Euler's brick*, Delta Epsilon, McGill Undergraduate Mathematics Journal **1** (2006), 30–33.
37. Knill O., *Hunting for Perfect Euler Bricks*, Harvard College Math. Review **2** (2008), no. 2, 102; see also <http://www.math.harvard.edu/~knill/various/eulercuboid/index.html>.
38. Sloan N. J. A., *Sequences A031173, A031174, and A031175*, On-line encyclopedia of integer sequences, OEIS Foundation Inc., Portland, USA.
39. Stoll M., Testa D., *The surface parametrizing cuboids*, e-print [arXiv:1009.0388](http://arXiv.org/abs/1009.0388) in Electronic Archive <http://arXiv.org>.
40. Sharipov R. A., *A note on a perfect Euler cuboid.*, e-print [arXiv:1104.1716](http://arXiv.org/abs/1104.1716) in Electronic Archive <http://arXiv.org>.
41. Sharipov R. A., *Perfect cuboids and irreducible polynomials*, Ufa Mathematical Journal **4**, (2012), no. 1, 153–160; see also e-print [arXiv:1108.5348](http://arXiv.org/abs/1108.5348) in Electronic Archive <http://arXiv.org>.
42. Sharipov R. A., *A note on the first cuboid conjecture*, e-print [arXiv:1109.2534](http://arXiv.org/abs/1109.2534) in Electronic Archive <http://arXiv.org>.
43. Sharipov R. A., *A note on the second cuboid conjecture. Part I*, e-print [arXiv:1201.1229](http://arXiv.org/abs/1201.1229) in Electronic Archive <http://arXiv.org>.
44. Sharipov R. A., *A note on the third cuboid conjecture. Part I*, e-print [arXiv:1203.2567](http://arXiv.org/abs/1203.2567) in Electronic Archive <http://arXiv.org>.
45. Sharipov R. A., *Perfect cuboids and multisymmetric polynomials*, e-print [arXiv:1205.3135](http://arXiv.org/abs/1205.3135) in Electronic Archive <http://arXiv.org>.
46. Sharipov R. A., *On an ideal of multisymmetric polynomials associated with perfect cuboids*, e-print [arXiv:1206.6769](http://arXiv.org/abs/1206.6769) in Electronic Archive <http://arXiv.org>.
47. Sharipov R. A., *On the equivalence of cuboid equations and their factor equations*, e-print [arXiv:1207.2102](http://arXiv.org/abs/1207.2102) in Electronic Archive <http://arXiv.org>.
48. Sharipov R. A., *A biquadratic Diophantine equation associated with perfect cuboids*, e-print [arXiv:1207.4081](http://arXiv.org/abs/1207.4081) in Electronic Archive <http://arXiv.org>.
49. Ramsden J. R., *A general rational solution of an equation associated with perfect cuboids*, e-print [arXiv:1207.5339](http://arXiv.org/abs/1207.5339) in Electronic Archive <http://arXiv.org>.
50. Ramsden J. R., Sharipov R. A., *Inverse problems associated with perfect cuboids*, e-print [arXiv:1207.6764](http://arXiv.org/abs/1207.6764) in Electronic Archive <http://arXiv.org>.

CSR PLC, CAMBRIDGE BUSINESS PARK, CAMBRIDGE, CB4 0WZ, UK
 E-mail address: jhnrsdn@yahoo.co.uk

BASHKIR STATE UNIVERSITY, 32 ZAKI VALIDI STREET, 450074 UFA, RUSSIA
 E-mail address: r-sharipov@mail.ru