

**A NOTE ON RATIONAL AND ELLIPTIC CURVES
ASSOCIATED WITH THE CUBOID FACTOR EQUATIONS.**

RUSLAN SHARIPOV

ABSTRACT. A rational perfect cuboid is a rectangular parallelepiped whose edges and face diagonals are given by rational numbers and whose space diagonal is equal to unity. It is described by a system of four equations with respect to six variables. The cuboid factor equations were derived from these four equations by symmetrization procedure. They constitute a system of eight polynomial equations, which has been solved parametrically. In the present paper its parametric solution is expressed through intersections or rational and elliptic curves arranged into parametric families.

1. INTRODUCTION.

Finding a rational perfect cuboid is equivalent to finding a perfect cuboid with all integer edges and diagonals, which is an old unsolved problem (see [1–44]). Here are the equations describing perfect cuboids:

$$\begin{aligned} p_0 &= x_1^2 + x_2^2 + x_3^2 - L^2 = 0, & p_1 &= x_2^2 + x_3^2 - d_1^2 = 0, \\ p_2 &= x_3^2 + x_1^2 - d_2^2 = 0, & p_3 &= x_1^2 + x_2^2 - d_3^2 = 0. \end{aligned} \quad (1.1)$$

The variables x_1, x_2, x_3 in (1.1) represent three edges of a cuboid, the variables d_1, d_2, d_3 are its face diagonals, and L is its space diagonal. In the case of a rational perfect cuboid we set $L = 1$. Using the polynomials p_0, p_1, p_2 , and p_3 from (1.1), we write the following eight polynomial equations:

$$\begin{aligned} p_0 &= 0, & \sum_{i=1}^3 p_i &= 0, \\ \sum_{i=1}^3 d_i p_i &= 0, & \sum_{i=1}^3 x_i p_i &= 0, \\ \sum_{i=1}^3 d_i^2 p_i &= 0, & \sum_{i=1}^3 x_i^2 p_i &= 0, \\ \sum_{i=1}^3 x_i d_i p_i &= 0, & \sum_{i=1}^3 x_i^2 d_i^2 p_i &= 0. \end{aligned} \quad (1.2)$$

2000 *Mathematics Subject Classification.* 11D25, 11D72, 14E05, 14H45, 14H52.

The equations (1.2) are called the cuboid factor equations. They were derived from the original cuboid equations (1.1) as a result of a symmetry approach initiated in [45] (see also [46–48]).

It is easy to see that each solution of the original cuboid equations (1.1) is a solution for the factor equations (1.2). Generally speaking, the converse is not true. However, in [47] the following theorem was proved.

Theorem 1.1. *Each integer or rational solution of the cuboid factor equations (1.2) such that $x_1 > 0$, $x_2 > 0$, $x_3 > 0$, $d_1 > 0$, $d_2 > 0$, and $d_3 > 0$ is an integer or rational solution for the original cuboid equations (1.1).*

Due to Theorem 1.1 the cuboid factor equations (1.2) are equivalent to the equations (1.1) in studying perfect cuboids. But in this paper, saying a solution of the factor equations, we assume any integer or rational solution, i. e. even such that some of the inequalities $x_1 > 0$, $x_2 > 0$, $x_3 > 0$, $d_1 > 0$, $d_2 > 0$, $d_3 > 0$ or all of them are not fulfilled.

The left hand sides of the cuboid factor equations are multisymmetric polynomials in x_1 , x_2 , x_3 and d_1 , d_2 , d_3 , i. e. they are invariant with respect to the permutation group S_3 acting upon x_1 , x_2 , x_3 , d_1 , d_2 , d_3 , and L as follows:

$$\sigma(x_i) = x_{\sigma i}, \quad \sigma(d_i) = d_{\sigma i}, \quad \sigma(L) = L.$$

For the theory of multisymmetric polynomials the reader is referred to [49–69]. According to this theory, each multisymmetric polynomial of x_1 , x_2 , x_3 and d_1 , d_2 , d_3 is expressed through the following nine elementary multisymmetric polynomials:

$$\begin{aligned} x_1 + x_2 + x_3 &= E_{10}, \\ x_1 x_2 + x_2 x_3 + x_3 x_1 &= E_{20}, \\ x_1 x_2 x_3 &= E_{30}, \end{aligned} \tag{1.3}$$

$$\begin{aligned} d_1 + d_2 + d_3 &= E_{01}, \\ d_1 d_2 + d_2 d_3 + d_3 d_1 &= E_{02}, \\ d_1 d_2 d_3 &= E_{03}, \end{aligned} \tag{1.4}$$

$$\begin{aligned} x_1 x_2 d_3 + x_2 x_3 d_1 + x_3 x_1 d_2 &= E_{21}, \\ x_1 d_2 + d_1 x_2 + x_2 d_3 + d_2 x_3 + x_3 d_1 + d_3 x_1 &= E_{11}, \\ x_1 d_2 d_3 + x_2 d_3 d_1 + x_3 d_1 d_2 &= E_{12}. \end{aligned} \tag{1.5}$$

Expressing the left hand sides of the cuboid factor equations (1.2) through the polynomials (1.3), (1.4), and (1.5), one gets polynomial equations with respect to the variables E_{10} , E_{20} , E_{30} , E_{01} , E_{02} , E_{03} , E_{21} , E_{11} , E_{12} , and L (see (3.1) through (3.7) in [48]). These equations were complemented with fourteen identities expressing the mutual algebraic dependence of the elementary multisymmetric polynomials (1.3), (1.4), and (1.5) (see (3.8) in [48]). As a result a system of twenty two equations was obtained. In [48] this huge system of twenty two equations was reduced to the following single polynomial equation for E_{10} , E_{01} , E_{11} , and L :

$$(2 E_{11})^2 + (E_{01}^2 + L^2 - E_{10}^2)^2 - 8 E_{01}^2 L^2 = 0. \tag{1.6}$$

The other variables E_{20} , E_{30} , E_{02} , E_{03} , E_{21} , E_{12} are expressed as rational functions of E_{10} , E_{01} , E_{11} , and L (see formulas (4.1), (4.3), (5.1), (5.2), (4.6), (4.7) in [48]).

The equation (1.6) was solved by John Ramsden in [70]. In the case of a rational perfect cuboid, where $L = 1$, omitting some inessential special cases, the general solution of the equation (1.6) is given by the formulas

$$E_{11} = -\frac{b(c^2 + 2 - 4c)}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}, \quad (1.7)$$

$$E_{10} = -\frac{b^2 c^2 + 2b^2 - 3b^2 c - c}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}, \quad (1.8)$$

$$E_{01} = -\frac{b(c^2 + 2 - 2c)}{b^2 c^2 + 2b^2 - 3b^2 c + c - b c^2 + 2b}. \quad (1.9)$$

Below are the formulas for E_{12} , E_{21} , E_{03} , E_{30} , E_{02} , E_{20} in (1.3), (1.4), and (1.5):

$$\begin{aligned} E_{12} = & (16b^6 + 32b^5 - 6c^5 b^2 + 2c^5 b - 62b^5 c^6 + 62b^6 c^6 + 16b^4 - \\ & - 180b^6 c^5 - c^7 b^3 + 18b^5 c^7 - 12b^6 c^7 - 2b^5 c^8 + b^6 c^8 + 248b^5 c^2 + \\ & + 248b^6 c^2 - 96b^6 c + 321b^6 c^4 - 180b^5 c^3 - 144b^5 c - 360b^6 c^3 + \\ & + b^4 c^8 + 8b^4 c^6 - 6b^4 c^7 + 18b^4 c^5 + 7b^3 c^6 + 90b^5 c^5 - 14b^3 c^5 + \\ & + 17b^2 c^4 + 32b^4 c^2 + 28b^3 c^3 - 28b^3 c^2 - 4b c^3 + 8b^3 c - 57b^4 c^4 + \\ & + 36b^4 c^3 - 12b^2 c^3 - 48b^4 c - c^4)(b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ & - 12b^2 c + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}, \end{aligned} \quad (1.10)$$

$$\begin{aligned} E_{21} = & \frac{b}{2} (5c^6 b - 2c^6 b^2 + 52c^5 b^2 - 16c^5 b - 2c^7 b^2 + 2b^4 c^8 - \\ & - 26b^4 c^7 - 426b^4 c^5 - 61b^3 c^6 + 100b^3 c^5 + 14c^7 b^3 - c^8 b^3 - 20b c^2 - \\ & - 8b^2 c^2 - 16b^2 c - 128b^2 c^4 - 200b^3 c^3 + 244b^3 c^2 + 32b c^3 + \\ & + 768b^4 c^4 - 852b^4 c^3 + 568b^4 c^2 + 104b^2 c^3 - 208b^4 c + 8c^4 + \\ & + 16b^3 - 112b^3 c + 142b^4 c^6 + 32b^4 - 2c^5)(b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ & - 12b^2 c - 4c^3 + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (bc - c - 2b)^{-2}, \end{aligned} \quad (1.11)$$

$$\begin{aligned} E_{03} = & \frac{b}{2} (b^2 c^4 - 5b^2 c^3 + 10b^2 c^2 - 10b^2 c + 4b^2 + 2bc + 2c^2 - \\ & - b c^3)(2b^2 c^4 - 12b^2 c^3 + 26b^2 c^2 - 24b^2 c + 8b^2 - c^4 b + 3b c^3 - \\ & - 6bc + 4b + c^3 - 2c^2 + 2c)(b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - \\ & - 12b^2 c + 4b^2 + c^2)^{-1} (bc - 1 - b)^{-2} (-c + bc - 2b)^{-2}, \end{aligned} \quad (1.12)$$

$$\begin{aligned} E_{30} = & c b^2 (1 - c)(c - 2)(b c^2 - 4bc + 2 + 4b)(2b c^2 - c^2 - 4bc + \\ & + 2b)(b^2 c^4 - 6b^2 c^3 + 13b^2 c^2 - 12b^2 c + 4b^2 + c^2)^{-1} \times \\ & \times (bc - 1 - b)^{-2} (-c + bc - 2b)^{-2}, \end{aligned} \quad (1.13)$$

$$\begin{aligned}
E_{02} = & \frac{1}{2} (28b^2c^2 - 16b^2c - 2c^2 - 4b^2 - b^2c^4 + 4b^3c^4 - 12b^3c^3 + \\
& + 4bc^3 + 24b^3c - 8bc - 2b^4c^4 + 12b^4c^3 - 26b^4c^2 - 8b^2c^3 + \\
& + 24b^4c - 16b^3 - 8b^4)(bc - 1 - b)^{-2}(bc - c - 2b)^{-2},
\end{aligned} \tag{1.14}$$

$$\begin{aligned}
E_{20} = & \frac{b}{2} (bc^2 - 2c - 2b)(2bc^2 - c^2 - 6bc + 2 + 4b) \times \\
& \times (bc - 1 - b)^{-2}(bc - c - 2b)^{-2}.
\end{aligned} \tag{1.15}$$

The formulas (1.10), (1.11), (1.12), (1.13), (1.14), (1.15) were derived in [71] by substituting the formulas (1.7), (1.8), and (1.9) along with $L = 1$ into the corresponding formulas from [48].

Thus, the right hand sides of the equalities (1.3), (1.4), and (1.5) turned out to be expressed through two arbitrary rational parameters b and c . The next step was to resolve these equalities with respect to $x_1, x_2, x_3, d_1, d_2, d_3$. For this purpose in [71] the following two cubic equations were written:

$$x^3 - E_{10}x^2 + E_{20}x - E_{30} = 0, \tag{1.16}$$

$$d^3 - E_{01}d^2 + E_{02}d - E_{03} = 0. \tag{1.17}$$

Note that the left hand sides of the equalities (1.3) are regular symmetric polynomials of the variables x_1, x_2, x_3 (see [72]). Similarly, the left hand sides of the equalities (1.4) are regular symmetric polynomials of the variables d_1, d_2, d_3 . For this reason x_1, x_2, x_3 can be found as roots of the cubic equation (1.16). Similarly, d_1, d_2, d_3 are roots of the second cubic equation (1.17). Relying on these facts, in [71] the following two inverse problems were formulated.

Problem 1.1. *Find all pairs of rational numbers b and c for which the cubic equations (1.16) and (1.17) with the coefficients given by the formulas (1.8), (1.9), (1.12), (1.13), (1.14), (1.15) possess positive rational roots $x_1, x_2, x_3, d_1, d_2, d_3$ obeying the auxiliary polynomial equations (1.5) whose right hand sides are given by the formulas (1.7), (1.10), and (1.11).*

Problem 1.2. *Find at least one pair of rational numbers b and c for which the cubic equations (1.16) and (1.17) with the coefficients given by the formulas (1.8), (1.9), (1.12), (1.13), (1.14), (1.15) possess positive rational roots $x_1, x_2, x_3, d_1, d_2, d_3$ obeying the auxiliary polynomial equations (1.5) whose right hand sides are given by the formulas (1.7), (1.10), and (1.11).*

Due to Theorem 1.1 the inverse problems 1.1 and 1.2 are equivalent to finding all rational perfect cuboids and to finding at least one rational perfect cuboid respectively. Singularities of the inverse problems 1.1 and 1.2 due to the denominators in the formulas (1.7) through (1.15) were studied in [73]. Some special cases where the equations (1.3), (1.4), (1.5) are solvable with respect to the cuboid variables x_1, x_2, x_3 and d_1, d_2, d_3 were found in [74]. However, none of these special cases have produced a perfect cuboid since the inequalities

$$\begin{aligned}
x_1 > 0, & & x_2 > 0, & & x_3 > 0, \\
d_1 > 0, & & d_2 > 0, & & d_3 > 0
\end{aligned} \tag{1.18}$$

required for solving the problems 1.1 and 1.2 are not fulfilled in these special cases.

Again, neglecting the inequalities (1.18), an approach to solving the equations (1.3), (1.4), (1.5) was found in [75]. It exploits the following lemma.

Lemma 1.1. *A reduced cubic equation $y^3 + y^2 + D = 0$ has three rational roots if and only if there is a rational number w satisfying the sextic equation*

$$D(w^2 + 3)^3 + 4(w - 1)^2(1 + w)^2 = 0. \quad (1.19)$$

In this case the roots of the cubic equation $y^3 + y^2 + D = 0$ are given by the formulas

$$y_1 = -\frac{2(w+1)}{w^2+3}, \quad y_2 = \frac{2(w-1)}{w^2+3}, \quad y_3 = \frac{1-w^2}{w^2+3}.$$

The idea of Lemma 1.1 belongs to John Ramsden. Its detailed proof is given in [75], where this lemma was applied to (1.16) and (1.17) (see Lemma 2.1 in [75]). As a result two sextic equations of the form (1.19) were derived:

$$D_1(w_1^2 + 3)^3 + 4(w_1 - 1)^2(1 + w_1)^2 = 0, \quad (1.20)$$

$$D_2(w_2^2 + 3)^3 + 4(w_2 - 1)^2(1 + w_2)^2 = 0. \quad (1.21)$$

The D -parameters D_1 and D_2 of the sextic equations (1.20) and (1.21) depend on the same two rational numbers b and c as E_{11} , E_{10} , E_{01} , E_{12} , E_{21} , E_{03} , E_{30} , E_{02} , E_{20} in the formulas (1.7) through (1.15). Therefore the equations (1.20) and (1.21) define two algebraic surfaces or, which is equivalent, two algebraic functions $w_1(b, c)$ and $w_2(b, c)$. Here are the explicit formulas for the D -parameters D_1 and D_2 in the sextic equations (1.20) and (1.21):

$$\begin{aligned} D_1 = & -\frac{2}{27}(7812b^4c^4 - 216b^2c^4 - 52b^2c^3 + 1764b^3c^4 - 1200b^4c^3 - \\ & - 1848b^4c^2 + 720b^4c - 36c^4b - 1512b^3c^3 - 36c^8b^3 + 288b^3c^2 - \\ & - 108c^6b^2 + 380c^5b^2 + 378c^7b^3 - 231c^8b^4 - 300c^7b^4 + 3906c^6b^4 - \\ & - 13c^7b^2 - 8904c^5b^4 - 882c^6b^3 + 18c^6b - 1319b^6c^8 + 20952b^5c^3 - \\ & - 11952b^5c^2 + 2592b^5c - 48372b^6c^4 + 31620b^6c^3 - 10552b^6c^2 + \\ & + 816b^6c + 1494b^5c^8 - 5238b^5c^7 - 4c^5 + 7905b^6c^7 - 24186b^6c^6 + \\ & + 288b^6 + 43740b^6c^5 + 7686b^5c^6 + 576b^7 + 128b^8 - 15372b^5c^4 - \\ & - 1080b^7c^8 - 3546b^7c^6 + 51c^9b^6 + 400b^8c^8 - 162c^9b^5 + 8640b^7c^2 - \\ & - 3456b^7c + 2808b^7c^7 - 1560b^8c^7 + 3940b^8c^6 + 216c^9b^7 - 960b^8c - \\ & - 6240b^8c^3 + 9c^{10}b^6 + 7880b^8c^4 + 4c^{10}b^8 - 6732b^8c^5 + 45c^9b^4 + \\ & + 3200b^8c^2 - 11232b^7c^3 + 7092b^7c^4 - 18c^{10}b^7 - 60c^9b^8)^2(2c^2 + \\ & + 2b^4c^4 - 12b^4c^3 + 26b^4c^2 - 24b^4c + 8b^4 - 6b^3c^4 + 18b^3c^3 - \\ & - 36b^3c + 24b^3 + 3b^2c^4 + 8b^2c^3 - 36b^2c^2 + 16b^2c + 12b^2 - 6bc^3 + \\ & + 12bc)^{-3}(b^2c^4 - 6b^2c^{-3} + 13b^2c^2 - 12b^2c + 4b^2 + c^2)^{-2}, \end{aligned} \quad (1.22)$$

$$\begin{aligned}
D_2 = & -\frac{2b^2}{27} (832b^2c^2 - 1440b^2c^4 - 840b^2c^3 + 4788b^3c^4 + 396bc^3 + \\
& + 720b^3c + 808b^4c^4 + 3032b^4c^3 - 2576b^4c^2 - 96b^4c + 448b^4 - \\
& - 504c^4b - 4176b^3c^3 - 9c^8b^3 + 72b^3c^2 - 720c^6b^2 + 2288c^5b^2 + \\
& + 1044c^7b^3 - 322c^8b^4 + 758c^7b^4 + 404c^6b^4 - 210c^7b^2 - 2464c^5b^4 - \\
& - 2394c^6b^3 + 72c^4 + 252c^6b + 3168b^6c^8 + 441c^9b^5 - 7056b^5c + \\
& + 57960b^6c^4 - 47232b^6c^3 + 25344b^6c^2 - 8064b^6c - 1809b^5c^8 + \\
& + 14472b^5c^2 + 3951b^5c^7 - 72c^5 + 36c^6 - 11808b^6c^7 + 1440b^5 + \\
& + 28980b^6c^6 - 49032b^6c^5 - 4410b^5c^6 + 8820b^5c^4 - 15804b^5c^3 + \\
& + 1152b^6 - 504c^9b^6 - 45c^9b^3 - 6c^9b^4 + 104c^8b^2 + 36c^{10}b^6 + \\
& + 14c^{10}b^4 - 45c^{10}b^5 - 99c^7b)^2 (6b^4c^4 - 36b^4c^3 + 78b^4c^2 - 72b^4c + \\
& + 24b^4 - 12b^3c^4 + 36b^3c^3 - 72b^3c + 48b^3 + 5b^2c^4 + 16b^2c^3 - \\
& - 68b^2c^2 + 32b^2c + 20b^2 - 12bc^3 + 24bc + 6c^2)^{-3} (b^2c^4 - 6b^2c^3 + \\
& + 13b^2c^2 - 12b^2c + 4b^2 + c^2)^{-2}.
\end{aligned} \tag{1.23}$$

The main result of [75] is expressed by the following two theorems.

Theorem 1.2. *Each rational point of the algebraic surface (1.20), except for points belonging to some algebraic subvariety of this surface, determine six rational numbers x_1, x_2, x_3 and d_1, d_2, d_3 obeying the cuboid factor equations (1.2) as well as the equations (1.3), (1.4), (1.5) whose right hand sides are given by the formulas (1.7) through (1.15).*

Theorem 1.3. *Each rational point of the algebraic surface (1.21), except for points belonging to some algebraic subvariety of this surface, determine six rational numbers x_1, x_2, x_3 and d_1, d_2, d_3 obeying the cuboid factor equations (1.2) as well as the equations (1.3), (1.4), (1.5) whose right hand sides are given by the formulas (1.7) through (1.15).*

The rational numbers x_1, x_2, x_3 and d_1, d_2, d_3 in Theorem 1.2 are given by explicit formulas expressing them as rational functions of b, c and w_1 :

$$x_i = x_i(b, c, w_1), \quad d_i = d_i(b, c, w_1). \tag{1.24}$$

But the formulas for $x_i(b, c, w_1)$ and $d_i(b, c, w_1)$ in (1.24) are very huge. For this reason they are not presented here. They are given in a machine readable form in the ancillary file [Solution.1.txt](#) to [76]. As for the exceptional subvariety mentioned in Theorem 1.2, it is determined by the denominator of (1.22) and by the denominators in those huge formulas for x_1, x_2, x_3 and d_1, d_2, d_3 .

The rational numbers x_1, x_2, x_3 and d_1, d_2, d_3 in Theorem 1.3 are also given by explicit formulas expressing them as rational functions of b, c and w_2 :

$$x_i = x_i(b, c, w_2), \quad d_i = d_i(b, c, w_2). \tag{1.25}$$

These formulas for $x_i(b, c, w_2)$ and $d_i(b, c, w_2)$ in (1.25) are also very huge. For this reason we do not present them here. They are given in a machine readable

form in the ancillary file [Solution_2.txt](#) to [76]. The exceptional subvariety in Theorem 1.3 is determined by the denominator of (1.23) and by the denominators in those huge formulas for x_1, x_2, x_3 and d_1, d_2, d_3 .

The algebraic surfaces (1.20) and (1.21) are not independent. They were studied in [76]. As it was shown in [76], these two algebraic surfaces are birationally equivalent. This birational equivalence is established by two rational functions

$$w_2 = w_2(b, c, w_1), \quad w_1 = w_1(b, c, w_2). \quad (1.26)$$

The formulas (1.26) should be treated modulo the equations (1.20) and (1.21) respectively. Then they produce two mutually inverse transformations.

The functions (1.24) and (1.25) are also not independent. They are related to each other through the birational transformations (1.26):

$$\begin{aligned} x_i(b, c, w_1) &= x_i(b, c, w_2(b, c, w_1)), \\ d_i(b, c, w_1) &= d_i(b, c, w_2(b, c, w_1)), \end{aligned} \quad (1.27)$$

$$\begin{aligned} x_i(b, c, w_2) &= x_i(b, c, w_1(b, c, w_2)), \\ d_i(b, c, w_2) &= d_i(b, c, w_1(b, c, w_2)). \end{aligned} \quad (1.28)$$

The explicit formulas for the rational functions $w_2(b, c, w_1)$ and $w_1(b, c, w_2)$ in (1.26) are very huge. They are not presented here, but they are given in a machine readable form in the ancillary file [Conversion_formulas.txt](#) to [76].

Due to (1.26), (1.27), and (1.28) the algebraic surfaces given by the sextic equations (1.20) and (1.21) are closely related to each other. Below we study both of them. Since $D_1 = D_1(b, c)$ and $D_2 = D_2(b, c)$ in (1.20) and (1.21), they are surfaces in \mathbb{R}^3 . The main goal of the present paper is to embed these surfaces into \mathbb{R}^4 and show that each point of any one of them lies in the intersection of some genus zero curve and some genus one curve specific to this point.

Note that some genus one curves associated with perfect cuboids were considered in [77]. However, they correspond to some very special solutions of the cubic equations (1.16) and (1.17). Genus one curves considered in this paper cover the general case in the theory of the cuboid factor equations.

2. THE STRUCTURE OF D_1 AND D_2 AND THE α -PARAMETERS.

Let's consider the formulas (1.22) and (1.23) for D_1 and D_2 . Looking at these formulas, one can detect the following structure of the expressions for D_1 and D_2 :

$$D_1 = -\frac{(P_1)^2}{(Q_1)^3}, \quad D_2 = -\frac{(P_2)^2}{(Q_2)^3}. \quad (2.1)$$

The denominators Q_1 and Q_2 in (2.1) are given by the formulas

$$\begin{aligned} Q_1 &= \frac{3}{2} (2c^2 + 2b^4c^4 - 12b^4c^3 + 26b^4c^2 - 24b^4c + 8b^4 - \\ &\quad - 6b^3c^4 + 18b^3c^3 - 36b^3c + 24b^3 + 3b^2c^4 + 8b^2c^3 - \\ &\quad - 36b^2c^2 + 16b^2c + 12b^2 - 6bc^3 + 12bc), \end{aligned} \quad (2.2)$$

$$\begin{aligned}
Q_2 = \frac{3}{2} & (6b^4c^4 - 36b^4c^3 + 78b^4c^2 - 72b^4c + 24b^4 - 12b^3c^4 + \\
& + 36b^3c^3 - 72b^3c + 48b^3 + 5b^2c^4 + 16b^2c^3 - 68b^2c^2 + \\
& + 32b^2c + 20b^2 - 12bc^3 + 24bc + 6c^2). \tag{2.3}
\end{aligned}$$

The numerators P_1 and P_2 in (2.1) are given by similar formulas:

$$\begin{aligned}
P_1 = \frac{1}{2} & (7812b^4c^4 - 216b^2c^4 - 52b^2c^3 + 1764b^3c^4 - 1200b^4c^3 - \\
& - 1848b^4c^2 + 720b^4c - 36c^4b - 1512b^3c^3 - 36c^8b^3 + 288b^3c^2 - \\
& - 108c^6b^2 + 380c^5b^2 + 378c^7b^3 - 231c^8b^4 - 300c^7b^4 + 3906c^6b^4 - \\
& - 13c^7b^2 - 8904c^5b^4 - 882c^6b^3 + 18c^6b - 1319b^6c^8 + 20952b^5c^3 - \\
& - 11952b^5c^2 + 2592b^5c - 48372b^6c^4 + 31620b^6c^3 - 10552b^6c^2 + \\
& + 816b^6c + 1494b^5c^8 - 5238b^5c^7 - 4c^5 + 7905b^6c^7 - 24186b^6c^6 + \\
& + 288b^6 + 43740b^6c^5 + 7686b^5c^6 + 576b^7 + 128b^8 - 15372b^5c^4 - \\
& - 1080b^7c^8 - 3546b^7c^6 + 51c^9b^6 + 400b^8c^8 - 162c^9b^5 + 8640b^7c^2 - \\
& - 3456b^7c + 2808b^7c^7 - 1560b^8c^7 + 3940b^8c^6 + 216c^9b^7 - 960b^8c - \\
& - 6240b^8c^3 + 9c^{10}b^6 + 7880b^8c^4 + 4c^{10}b^8 - 6732b^8c^5 + 45c^9b^4 + \\
& + 3200b^8c^2 - 11232b^7c^3 + 7092b^7c^4 - 18c^{10}b^7 - 60c^9b^8) \times \\
& \times (b^2c^4 - 6b^2c^{-3} + 13b^2c^2 - 12b^2c + 4b^2 + c^2)^{-1}, \tag{2.4}
\end{aligned}$$

$$\begin{aligned}
P_2 = \frac{b}{2} & (832b^2c^2 - 1440b^2c^4 - 840b^2c^3 + 4788b^3c^4 + 396bc^3 + \\
& + 720b^3c + 808b^4c^4 + 3032b^4c^3 - 2576b^4c^2 - 96b^4c + 448b^4 - \\
& - 504c^4b - 4176b^3c^3 - 9c^8b^3 + 72b^3c^2 - 720c^6b^2 + 2288c^5b^2 + \\
& + 1044c^7b^3 - 322c^8b^4 + 758c^7b^4 + 404c^6b^4 - 210c^7b^2 - 2464c^5b^4 - \\
& - 2394c^6b^3 + 72c^4 + 252c^6b + 3168b^6c^8 + 441c^9b^5 - 7056b^5c + \\
& + 57960b^6c^4 - 47232b^6c^3 + 25344b^6c^2 - 8064b^6c - 1809b^5c^8 + \\
& + 14472b^5c^2 + 3951b^5c^7 - 72c^5 + 36c^6 - 11808b^6c^7 + 1440b^5 + \\
& + 28980b^6c^6 - 49032b^6c^5 - 4410b^5c^6 + 8820b^5c^4 - 15804b^5c^3 + \\
& + 1152b^6 - 504c^9b^6 - 45c^9b^3 - 6c^9b^4 + 104c^8b^2 + 36c^{10}b^6 + \\
& + 14c^{10}b^4 - 45c^{10}b^5 - 99c^7b) (b^2c^4 - 6b^2c^3 + \\
& + 13b^2c^2 - 12b^2c + 4b^2 + c^2)^{-2}. \tag{2.5}
\end{aligned}$$

Due to (2.1) and the formulas (2.2), (2.3), (2.4), (2.5) we can write the sextic equations (1.20) and (1.21) in the following way:

$$\left(\frac{w_1^2 + 3}{Q_1} \right)^3 = \left(\frac{2(w_1^2 - 1)}{P_1} \right)^2, \quad \left(\frac{w_2^2 + 3}{Q_2} \right)^3 = \left(\frac{2(w_2^2 - 1)}{P_2} \right)^2. \tag{2.6}$$

Lemma 2.1. *If x and y are two rational numbers obeying the equality $x^3 = y^2$, then there is a third rational number α such that $x = \alpha^2$ and $y = \alpha^3$.*

Proof. Each rational number is presented as an irreducible ratio of two integer numbers. Expanding these numbers into prime factors we find that each rational number is a product of distinct prime numbers to some integer powers which can be either positive or negative. For x and y this fact yields

$$x = \pm p_1^{\beta_1} \cdot \dots \cdot p_m^{\beta_m}, \quad y = \pm q_1^{\gamma_1} \cdot \dots \cdot q_n^{\gamma_n}. \quad (2.7)$$

Substituting (2.7) into the equality $x^3 = y^2$, we find that $x > 0$, $m = n$ and the prime factors p_1, \dots, p_m should coincide with the prime factors q_1, \dots, q_n up to some permutation. Performing this permutation upon the prime factors q_1, \dots, q_n , for the exponents β_1, \dots, β_m and $\gamma_1, \dots, \gamma_n$ in (2.7) we derive

$$3\beta_i = 2\gamma_i, \text{ where } i = 1, \dots, n. \quad (2.8)$$

Due to (2.8) there are some unique integer numbers $\alpha_1, \dots, \alpha_n$ such that

$$\beta_i = 2\alpha_i, \quad \gamma_i = 3\alpha_i. \quad (2.9)$$

Using the numbers $\alpha_1, \dots, \alpha_n$, we define the number α by setting

$$\alpha = \pm p_1^{\alpha_1} \cdot \dots \cdot p_n^{\alpha_n}. \quad (2.10)$$

The sign of α in (2.10) is chosen to be coinciding with the sign of y in (2.7). Then from (2.9), using $x > 0$ and (2.7), we derive the required equalities $x = \alpha^2$ and $y = \alpha^3$. Lemma 2.1 is proved. \square

Now we can apply Lemma 2.1 to the equations (2.6). As a result we obtain the following two theorems whose proofs are obvious.

Theorem 2.1. *For each rational point (b, c, w_1) of the algebraic surface (1.20) in \mathbb{R}^3 , where $w_1 \neq \pm 1$, there is a rational number α_1 such that*

$$w_1^2 + 3 = Q_1 \alpha_1^2, \quad 2(w_1^2 - 1) = P_1 \alpha_1^3. \quad (2.11)$$

The numbers P_1 and Q_1 in (2.11) are given by the formulas (2.2) and (2.4).

Theorem 2.2. *For each rational point (b, c, w_2) of the algebraic surface (1.21) in \mathbb{R}^3 , where $w_2 \neq \pm 1$, there is a rational number α_2 such that*

$$w_2^2 + 3 = Q_2 \alpha_2^2, \quad 2(w_2^2 - 1) = P_2 \alpha_2^3. \quad (2.12)$$

The numbers P_2 and Q_2 in (2.12) are given by the formulas (2.3) and (2.5).

3. GENUS ZERO CURVES.

Let's consider the equations (2.11) separately. Since $Q_1 = Q_1(b, c)$, the first equation (2.11) defines a hypersurface in \mathbb{R}^4 . Similarly, $P_1 = P_1(b, c)$. Therefore the second equation (2.11) defines another hypersurface in \mathbb{R}^4 . Thus, we get an embedding of the algebraic surface (1.20) into \mathbb{R}^4 where this surface is presented as the intersection of two algebraic hypersurfaces. In a similar way, the equations (2.12) define an embedding of the algebraic surface (1.21) into \mathbb{R}^4 where this surface

is presented as the intersection of two algebraic hypersurfaces.

Now assume that the rational numbers b and c are chosen to be constants. Then P_1 and Q_1 in (2.11) are also two rational constants. Under this assumption the first equation (2.11) defines a genus zero curve in \mathbb{R}^2 , while the second equation defines a genus one curve. The same is true for the equations (2.12), provided b and c are considered as constants.

Definition 3.1. A genus zero algebraic curve is called a rational curve if it is birationally equivalent to a line (see [78]).

Theorem 3.1. An algebraic curve in \mathbb{R}^2 given by the quadratic equation

$$w^2 + 3 = Q \alpha^2, \quad (3.1)$$

where Q is a rational number, is birationally equivalent to a line over \mathbb{Q} if and only if it has at least one rational point (w_0, α_0) in \mathbb{R}^2 .

Proof. Necessity. Assume that the curve (3.1) is birationally equivalent to a line over \mathbb{Q} . Then there are two rational functions with rational coefficients

$$w = w(t), \quad \alpha = \alpha(t) \quad (3.2)$$

whose domain is some Zariski open subset $D \subseteq \mathbb{Q} \subset \mathbb{R}$ and such that they satisfy the equality (3.1) identically. Choosing some point $t_0 \in D$ and substituting $t = t_0$ into (3.2), we get two rational numbers $w_0 = w(t_0)$ and $\alpha_0 = \alpha(t_0)$ satisfying the equality (3.1). They constitute a rational point (w_0, α_0) of our curve (3.1).

Sufficiency. Assume that (w_0, α_0) is some rational point of the curve (3.1). Then the following equality holds for w_0 and α_0 :

$$w_0^2 + 3 = Q \alpha_0^2. \quad (3.3)$$

Subtracting (3.3) from (3.1), we derive

$$(w - w_0)(w + w_0) = Q(\alpha - \alpha_0)(\alpha + \alpha_0). \quad (3.4)$$

The equality (3.4) can be transformed to the following one:

$$\frac{w - w_0}{\alpha + \alpha_0} = Q \frac{\alpha - \alpha_0}{w + w_0}. \quad (3.5)$$

Using the equality (3.5), we introduce the parameter t by setting

$$\frac{\alpha - \alpha_0}{w + w_0} = t, \quad \frac{w - w_0}{\alpha + \alpha_0} = Q t. \quad (3.6)$$

The equations (3.6) are easily resolved with respect to w and α :

$$w = \frac{w_0 + 2Q\alpha_0 t + Qw_0 t^2}{1 - Qt^2}, \quad \alpha = \frac{\alpha_0 + 2w_0 t + Q\alpha_0 t^2}{1 - Qt^2}. \quad (3.7)$$

The functions (3.7) are that very rational functions $w(t)$ and $\alpha(t)$ which implement the birational equivalence of the curve (3.1) and a line. Theorem is proved. \square

Note that Q in (3.1) is a rational number. Therefore $Q = (M m^2)/(N n^2)$, where M and N are relatively prime square free integer numbers. Let (w_0, α_0) be a rational point of the algebraic curve (3.1). Then w_0 and $(\alpha_0 m)/(N n)$ are two rational numbers. We can express these numbers as

$$w_0 = \frac{X}{Z}, \quad \frac{\alpha_0 m}{N n} = \frac{Y}{Z}, \quad (3.8)$$

where X , Y , and Z are three integers. Applying $Q = (M m^2)/(N n^2)$ and (3.8) to (3.1), we derive the following Diophantine equation:

$$X^2 - M N Y^2 + 3 Z^2 = 0. \quad (3.9)$$

The Diophantine equation (3.9) is an instance of the Legendre equation (see [79], [80], and [81]). There is a criterion for the equation (3.9) to have a solution. This criterion can be derived from the following Legendre theorem.

Theorem 3.2 (Legendre). *A quadratic Diophantine equation with square free and pairwise relatively prime coefficients*

$$A X^2 + B Y^2 + C Z^2 = 0 \quad (3.10)$$

has a nonzero solution if and only if its coefficients are not all of the same sign and if $-BC$, $-CA$, and $-AB$ are squares modulo A , B , and C respectively.

Due to (3.8) the numbers M and N in (3.9) are relatively prime and square free. Therefore their product $M N$ is also square free. The equation (3.9) is a special instance of the equation (3.10). There are special solvability criteria for it (see [80]).

Theorem 3.3. *If $M N$ is nonzero modulo 3, then the Diophantine equation (3.9) has a nonzero solution if and only if the following three conditions are fulfilled:*

- 1) $M N > 0$, 2) -3 is a square modulo $M N$, 3) $M N$ is a square modulo 3.

Theorem 3.4. *If $M N$ is zero modulo 3, then the Diophantine equation (3.9) has a nonzero solution if and only if the following three conditions are fulfilled:*

- 1) $M N > 0$, 2) -3 is a square modulo $M N$, 3) $\frac{M N}{3}$ is a square modulo 3.

Applying Theorems 3.3 and 3.4, we can study the rationality of genus zero curves from (2.11) and (2.12) for particular numeric values of b and c :

$$w_1^2 + 3 = Q_1 \alpha_1^2, \quad w_2^2 + 3 = Q_2 \alpha_2^2. \quad (3.11)$$

The results of such numeric studies for the curves (3.11) are presented in the ancillary file **Quadratic_curves.txt** attached to this arXiv submission.

Looking through the numeric output file **Quadratic_curves.txt**, we see that some genus zero curves are rational over \mathbb{Q} . Others are not rational. So, both types of curves arise in the theory of cuboid factor equations.

4. GENUS ONE CURVES.

Genus one algebraic curves in (2.11) and (2.12) are more regular than genus zero curves (3.11). They are given by the following cubic equations:

$$2(w_1^2 - 1) = P_1 \alpha_1^3, \quad 2(w_2^2 - 1) = P_2 \alpha_2^3. \quad (4.1)$$

Definition 4.1. A genus one algebraic curve is called an elliptic curve if it has at least one rational point either finite or at infinity (see [82]).

The parameters P_1 and P_2 in (4.1) are given by the formulas (2.4) and (2.5). But despite the values of these parameters, both curves (4.1) have rational points

$$(w_1, \alpha_1) = (\pm 1, 0), \quad (w_2, \alpha_2) = (\pm 1, 0). \quad (4.2)$$

The points (4.2) are exceptional in the sense of Theorems 2.1 and 2.2. Nevertheless, they are sufficient to say that both curves (4.1) are elliptic curves.

5. CONCLUDING REMARKS.

Summarizing the results of Section 3 and Section 4 above, we conclude that each rational point of the algebraic surfaces (1.20) and (1.21) is produced through the intersection of some elliptic curve and some rational curve. This fact could be used for to apply powerful tools of the modern theory of elliptic curves in studying the equations (1.20) and (1.21) and thus in studying perfect cuboids.

REFERENCES

1. *Euler brick*, Wikipedia, Wikimedia Foundation Inc., San Francisco, USA.
2. Halcke P., *Deliciae mathematicae oder mathematisches Sinnen-Confect*, N. Sauer, Hamburg, Germany, 1719.
3. Saunderson N., *Elements of algebra*, Vol. 2, Cambridge Univ. Press, Cambridge, 1740.
4. Euler L., *Vollständige Anleitung zur Algebra*, 3 Theile, Kaiserliche Akademie der Wissenschaften, St. Petersburg, 1770-1771.
5. Pocklington H. C., *Some Diophantine impossibilities*, Proc. Cambridge Phil. Soc. **17** (1912), 108–121.
6. Dickson L. E., *History of the theory of numbers*, Vol. 2: *Diophantine analysis*, Dover, New York, 2005.
7. Kraitchik M., *On certain rational cuboids*, Scripta Math. **11** (1945), 317–326.
8. Kraitchik M., *Théorie des Nombres*, Tome 3, *Analyse Diophantine et application aux cuboïdes rationnelles*, Gauthier-Villars, Paris, 1947.
9. Kraitchik M., *Sur les cuboïdes rationnelles*, Proc. Int. Congr. Math. **2** (1954), Amsterdam, 33–34.
10. Bromhead T. B., *On square sums of squares*, Math. Gazette **44** (1960), no. 349, 219–220.
11. Lal M., Blundon W. J., *Solutions of the Diophantine equations $x^2 + y^2 = l^2$, $y^2 + z^2 = m^2$, $z^2 + x^2 = n^2$* , Math. Comp. **20** (1966), 144–147.
12. Spohn W. G., *On the integral cuboid*, Amer. Math. Monthly **79** (1972), no. 1, 57–59.
13. Spohn W. G., *On the derived cuboid*, Canad. Math. Bull. **17** (1974), no. 4, 575–577.
14. Chein E. Z., *On the derived cuboid of an Eulerian triple*, Canad. Math. Bull. **20** (1977), no. 4, 509–510.
15. Leech J., *The rational cuboid revisited*, Amer. Math. Monthly **84** (1977), no. 7, 518–533; see also Erratum, Amer. Math. Monthly **85** (1978), 472.
16. Leech J., *Five tables relating to rational cuboids*, Math. Comp. **32** (1978), 657–659.
17. Spohn W. G., *Table of integral cuboids and their generators*, Math. Comp. **33** (1979), 428–429.
18. Lagrange J., *Sur le dérivé du cuboïde Eulérien*, Canad. Math. Bull. **22** (1979), no. 2, 239–241.

19. Leech J., *A remark on rational cuboids*, Canad. Math. Bull. **24** (1981), no. 3, 377–378.
20. Korec I., *Nonexistence of small perfect rational cuboid*, Acta Math. Univ. Comen. **42/43** (1983), 73–86.
21. Korec I., *Nonexistence of small perfect rational cuboid II*, Acta Math. Univ. Comen. **44/45** (1984), 39–48.
22. Wells D. G., *The Penguin dictionary of curious and interesting numbers*, Penguin publishers, London, 1986.
23. Bremner A., Guy R. K., *A dozen difficult Diophantine dilemmas*, Amer. Math. Monthly **95** (1988), no. 1, 31–36.
24. Bremner A., *The rational cuboid and a quartic surface*, Rocky Mountain J. Math. **18** (1988), no. 1, 105–121.
25. Colman W. J. A., *On certain semiperfect cuboids*, Fibonacci Quart. **26** (1988), no. 1, 54–57; see also *Some observations on the classical cuboid and its parametric solutions*, Fibonacci Quart. **26** (1988), no. 4, 338–343.
26. Korec I., *Lower bounds for perfect rational cuboids*, Math. Slovaca **42** (1992), no. 5, 565–582.
27. Guy R. K., *Is there a perfect cuboid? Four squares whose sums in pairs are square. Four squares whose differences are square*, Unsolved Problems in Number Theory, 2nd ed., Springer-Verlag, New York, 1994, pp. 173–181.
28. Rathbun R. L., Granlund T., *The integer cuboid table with body, edge, and face type of solutions*, Math. Comp. **62** (1994), 441–442.
29. Van Luijk R., *On perfect cuboids*, Doctoraalscriptie, Mathematisch Instituut, Universiteit Utrecht, Utrecht, 2000.
30. Rathbun R. L., Granlund T., *The classical rational cuboid table of Maurice Kraitchik*, Math. Comp. **62** (1994), 442–443.
31. Peterson B. E., Jordan J. H., *Integer hexahedra equivalent to perfect boxes*, Amer. Math. Monthly **102** (1995), no. 1, 41–45.
32. Rathbun R. L., *The rational cuboid table of Maurice Kraitchik*, e-print [math.HO/0111229](http://arXiv.org/math.HO/0111229) in Electronic Archive <http://arXiv.org>.
33. Hartshorne R., Van Luijk R., *Non-Euclidean Pythagorean triples, a problem of Euler, and rational points on $K3$ surfaces*, e-print [math.NT/0606700](http://arXiv.org/math.NT/0606700) in Electronic Archive <http://arXiv.org>.
34. Waldschmidt M., *Open diophantine problems*, e-print [math.NT/0312440](http://arXiv.org/math.NT/0312440) in Electronic Archive <http://arXiv.org>.
35. Ionascu E. J., Luca F., Stanica P., *Heron triangles with two fixed sides*, e-print [math.NT/0608185](http://arXiv.org/math.NT/0608185) in Electronic Archive <http://arXiv.org>.
36. Ortan A., Quenneville-Belair V., *Euler's brick*, Delta Epsilon, McGill Undergraduate Mathematics Journal **1** (2006), 30–33.
37. Knill O., *Hunting for Perfect Euler Bricks*, Harvard College Math. Review **2** (2008), no. 2, 102; see also <http://www.math.harvard.edu/~knill/various/eulercuboid/index.html>.
38. Sloan N. J. A., *Sequences A031173, A031174, and A031175*, On-line encyclopedia of integer sequences, OEIS Foundation Inc., Portland, USA.
39. Stoll M., Testa D., *The surface parametrizing cuboids*, e-print [arXiv:1009.0388](http://arXiv.org/abs/1009.0388) in Electronic Archive <http://arXiv.org>.
40. Sharipov R. A., *A note on a perfect Euler cuboid.*, e-print [arXiv:1104.1716](http://arXiv.org/abs/1104.1716) in Electronic Archive <http://arXiv.org>.
41. Sharipov R. A., *Perfect cuboids and irreducible polynomials*, Ufa Mathematical Journal **4**, (2012), no. 1, 153–160; see also e-print [arXiv:1108.5348](http://arXiv.org/abs/1108.5348) in Electronic Archive <http://arXiv.org>.
42. Sharipov R. A., *A note on the first cuboid conjecture*, e-print [arXiv:1109.2534](http://arXiv.org/abs/1109.2534) in Electronic Archive <http://arXiv.org>.
43. Sharipov R. A., *A note on the second cuboid conjecture. Part I*, e-print [arXiv:1201.1229](http://arXiv.org/abs/1201.1229) in Electronic Archive <http://arXiv.org>.
44. Sharipov R. A., *A note on the third cuboid conjecture. Part I*, e-print [arXiv:1203.2567](http://arXiv.org/abs/1203.2567) in Electronic Archive <http://arXiv.org>.
45. Sharipov R. A., *Perfect cuboids and multisymmetric polynomials*, e-print [arXiv:1205.3135](http://arXiv.org/abs/1205.3135) in Electronic Archive <http://arXiv.org>.
46. Sharipov R. A., *On an ideal of multisymmetric polynomials associated with perfect cuboids*, e-print [arXiv:1206.6769](http://arXiv.org/abs/1206.6769) in Electronic Archive <http://arXiv.org>.
47. Sharipov R. A., *On the equivalence of cuboid equations and their factor equations*, e-print [arXiv:1207.2102](http://arXiv.org/abs/1207.2102) in Electronic Archive <http://arXiv.org>.

48. Sharipov R. A., *A biquadratic Diophantine equation associated with perfect cuboids*, e-print arXiv:1207.4081 in Electronic Archive <http://arXiv.org>.
49. Shläfli L., *Über die Resultante eines systems mehrerer algebraischen Gleichungen*, Denkschr. Kaiserliche Acad. Wiss. Math.-Natur. Kl. **4** (1852); reprinted in «Gesammelte mathematische Abhandlungen», Band II (1953), Birkhäuser Verlag, 9–112.
50. Cayley A., *On the symmetric functions of the roots of certain systems of two equations*, Phil. Trans. Royal Soc. London **147** (1857), 717–726.
51. Junker F., *Über symmetrische Functionen von mehreren Veränderlichen*, Mathematische Annalen **43** (1893), 225–270.
52. McMahon P. A., *Memoir on symmetric functions of the roots of systems of equations*, Phil. Trans. Royal Soc. London **181** (1890), 481–536.
53. McMahon P. A., *Combinatory Analysis*. Vol. I and Vol. II, Cambridge Univ. Press, 1915–1916; see also Third ed., Chelsea Publishing Company, New York, 1984.
54. Noether E., *Der Endlichkeitssatz der Invarianten endlicher Gruppen*, Mathematische Annalen **77** (1915), 89–92.
55. Weyl H., *The classical groups*, Princeton Univ. Press, Princeton, 1939.
56. Macdonald I. G., *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1979.
57. Pedersen P., *Calculating multidimensional symmetric functions using Jacobi's formula*, Proceedings AAEECC 9, volume 539 of Springer Lecture Notes in Computer Science, Springer, 1991, pp. 304–317.
58. Milne P., *On the solutions of a set of polynomial equations*, Symbolic and numerical computation for artificial intelligence. Computational Mathematics and Applications (Donald B. R., Kapur D., Mundy J. L., eds.), Academic Press Ltd., London, 1992, pp. 89–101.
59. Dalbec J., *Geometry and combinatorics of Chow forms*, PhD thesis, Cornell University, 1995.
60. Richman D. R., *Explicit generators of the invariants of finite groups*, Advances in Math. **124** (1996), no. 1, 49–76.
61. Stepanov S. A., *On vector invariants of the symmetric group*, Diskretnaya Matematika **8** (1996), no. 2, 48–62.
62. Gonzalez-Vega L., Trujillo G., *Multivariate Sturm-Habicht sequences: real root counting on n -rectangles and triangles*, Revista Matemática Complutense **10** (1997), 119–130.
63. Stepanov S. A., *On vector invariants of symmetric groups*, Diskretnaya Matematika **11** (1999), no. 3, 4–14.
64. Dalbec J., *Multisymmetric functions*, Beiträge zur Algebra und Geom. **40** (1999), no. 1, 27–51.
65. Rosas M. H., *MacMahon symmetric functions, the partition lattice, and Young subgroups*, Journ. Combin. Theory **96 A** (2001), no. 2, 326–340.
66. Vaccarino F., *The ring of multisymmetric functions*, e-print math.RA/0205233 in Electronic Archive <http://arXiv.org>.
67. Briand E., *When is the algebra of multisymmetric polynomials generated by the elementary multisymmetric polynomials?*, Beiträge zur Algebra und Geom. **45** (2004), no. 2, 353–368.
68. Rota G.-C., Stein J. A., *A problem of Cayley from 1857 and how he could have solved it*, Linear Algebra and its Applications (special issue on determinants and the legacy of Sir Thomas Muir) **411** (2005), 167–253.
69. Briand E., Rosas M. H., *Milne's volume function and vector symmetric polynomials*, Journ. Symbolic Comput. **44** (2009), no. 5, 583–590.
70. Ramsden J. R., *A general rational solution of an equation associated with perfect cuboids*, e-print arXiv:1207.5339 in Electronic Archive <http://arXiv.org>.
71. Ramsden J. R., Sharipov R. A., *Inverse problems associated with perfect cuboids*, e-print arXiv:1207.6764 in Electronic Archive <http://arXiv.org>.
72. *Symmetric polynomial*, Wikipedia, Wikimedia Foundation Inc., San Francisco, USA.
73. Ramsden J. R., Sharipov R. A., *On singularities of the inverse problems associated with perfect cuboids*, e-print arXiv:1208.1859 in Electronic Archive <http://arXiv.org>.
74. Sharipov R. A., *On a pair of cubic equations associated with perfect cuboids*, e-print arXiv:1208.0308 in Electronic Archive <http://arXiv.org>.
75. Ramsden J. R., Sharipov R. A., *On two algebraic parametrizations for rational solutions of the cuboid equations*, e-print arXiv:1208.2587 in Electronic Archive <http://arXiv.org>.
76. Sharipov R. A., *A note on solutions of the cuboid factor equations*, e-print arXiv:1209.0723 in Electronic Archive <http://arXiv.org>.

77. Sharipov R. A., *On two elliptic curves associated with perfect cuboids*, e-print [arXiv:1208.1227](https://arxiv.org/abs/1208.1227) in Electronic Archive <http://arXiv.org>.
78. *Rational curves*, Wikipedia, Wikimedia Foundation Inc., San Francisco, USA.
79. *Legendre's equation*, Wikipedia, Wikimedia Foundation Inc., San Francisco, USA.
80. Aitken W., *Legendre's Theorem, Lagrange's Descent*, see on-line [Lecture Notes and Other Things of Interest](#), California State University, San Marcos, USA.
81. Cremona J. E., Rusin D., *Efficient solution of rational conics*, *Mathematics of Computation* **72** (2003), no. 243, 1417–1441.
82. Connel I., *Elliptic curve handbook*, McGill University, Montreal, 1999; see <http://www.math.mcgill.ca/connell>.

BASHKIR STATE UNIVERSITY, 32 ZAKI VALIDI STREET, 450074 UFA, RUSSIA
E-mail address: r-sharipov@mail.ru